


RESEARCH

Open Access



“The measures taken by the government overburdened the daily practice” – insights of the PRICOV-19 study on German general practitioners in times of COVID-19

Stefanie Stark^{1*} , Emmily Schaubroeck², Marie Kluge¹, Larissa Burggraf^{1,3}, Marco Roos⁴, Eve Borowski¹, Esther Van Poel², Sara Willems², Thomas Kühlein¹, Susann Hueber¹ and Felix Werner¹

Abstract

Background The international study PRICOV-19 aims to assess the impact of the COVID-19 pandemic on the organisation of primary health care. The German part focuses on German general practitioners during the second wave of the COVID-19 pandemic. This paper addresses the following research questions: (1) How were changes in tasks on primary care and patient treatment perceived by GPs?, (2) What was the role of GPs during the pandemic, and how was their wellbeing?, (3) How did GPs perceive health policy measures?, and, (4) What influenced the attitudes of GPs on health policy measures?

Methods This study pursues a multi-country cross-sectional design. Data collection took place throughout Germany from 01.02. to 28.02.2021 with a quantitative online questionnaire consisting of 53 items. The questionnaire was analysed through descriptive and inferential analyses using correlation and multiple regression models.

Results The response rate was 20.4% ($n = 349$). The respondents were mainly GPs (59.6%) in single practices (62.5%) with a mean work experience of 15 to 20 years. GPs experienced a change in their work and practice organisation (80.3%). They felt a high responsibility (70.6%) and found their work has become more meaningful to them (76%). They also saw a lack of political support (75.2%) and that the measures taken by the government overburdened the daily practice (66.4%). Not many GPs were at risk of being distressed (53.4%) but rated the health policies rather negatively (60%). The multiple regression showed, the more GPs were exposed to risk of distress, the worse they assessed the government's measures.

Conclusion GPs perceived their work as relevant and felt confident they could fulfil their tasks, but noticed that health policy initially hardly supported the outpatient sector. Health policies should increase their competence in relation to primary care, ensure its needs and consider an active inclusion of GPs in preparedness plans.

Keywords Primary health care, General practitioners, COVID-19, PRICOV-19, Quality of care, Health policy, Wellbeing

*Correspondence:

Stefanie Stark

Stefanie.Stark@uk-erlangen.de

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Due to the rapid spread and unforeseen course of the pandemic, the high contagiousness of the SARS-CoV-2 virus, and the high mortality rate, the COVID-19 pandemic presented itself as a worldwide crisis [1–5]. Health systems were heavily exposed to this pandemic and faced new challenges to medical-professional interactions [5, 6]. In Germany, this crisis revealed who or what was coined "system-relevant" [7, 8], i.e., necessary for the maintenance and continued existence of the healthcare system [6]. The first COVID-19 wave (March 2020 – May 2020) hit the German health system mostly unprepared [9]. The health policy, as well as the system, was not sufficiently prepared at that time. The general public first noticed the effects of the pandemic in the public health sector, however the first place where the effects became visible was the clinical sector, especially the intensive care units (ICU). Initially, research, media and particularly political focus were primarily on this sector in its fight against the virus [10–14]. Nevertheless, a comprehensive approach to managing the COVID-19 pandemic should also focus on primary care, as the outpatient sector was also caught unprepared by this pandemic: no or inadequate personal protective equipment (PPE), no instructions or measures for pandemic preparedness (PP), and initially hardly or insufficient support from health policies [15]. Albeit at this time, six out of seven COVID-19 patients in Germany were mainly treated in primary care practices first [2, 14, 16].

Accordingly, the prolonged COVID-19 crisis has triggered government and policy reactions that intended to protect the ICUs from being overburdened by maintaining the primary care sector functioning [17, 18]. Especially with the beginning of the second pandemic wave (September 2020 – February 2021) in Germany [9], primary care played an essential role in the care of infected patients and was put at the forefront of providing essential support in containing and diagnosing COVID-19 [19–21]. New regulations posed organisational, structural, and content-related challenges in German general practices [21–23]: e.g., the emergency service being used for COVID-19 testing, as well as the implementation of special infection consultation hours [24]. Primary care was set to be the first point of contact where decisions had to be taken on whether patients should be referred to the hospital for further treatment. Patients with less severe and uncomplicated symptoms were entirely cared for in primary care [25, 26]. Furthermore, preventive care, e.g., vaccinations, was widely accomplished by General practitioners (GPs) in their practices [19, 20, 27, 28]. In this pandemic, the primary care sector had to maintain basic

health care structure, especially if other parts of the healthcare system were in danger of overloading [4, 21, 29]. Therefore, German GPs functioned as gatekeepers who provided authorisation for access to hospital care by mitigating the risk of overburdened clinics and delayed specialist care [13, 30, 31]. Former studies have shown that primary care and its actors were under intense pressure and faced new challenges and difficulties during the COVID-19 pandemic [23, 26, 32] and has thus presented new and fast-changing tasks for GPs to maintain outpatient care [19, 21, 26, 33]. Furthermore, studies regarding previous pandemics demonstrated a negative impact on the GPs' perception of their role. Accordingly, previous research also showed that the COVID-19 pandemic, with its policy measures as well as its regulations, negatively affected the GPs' workload and, subsequently, their wellbeing [34, 35].

The international collaboration study *PRICOV-19 – Primary Health Care in times of COVID-19*¹ aims to assess the impact of the COVID-19 pandemic on the organisation of primary health care. The German part of this study focuses on the impact of the pandemic measures and health policy regulations on GPs' perceptions and wellbeing during the second wave of the COVID-19 pandemic in Germany (February 2021). It is assumed from the research aforementioned that pandemic-related changes due to associated political actions would manifest themselves in the role and tasks of the GPs, affecting their workload, as well as their mental wellbeing. Accordingly, this paper addresses the following research questions: (1) *How were changes in tasks on primary care and patient treatment perceived by GPs?*, (2) *What was the role of GPs during the second wave of the pandemic, and how was their wellbeing?*, (3) *How did GPs perceive the health policy measures?*, and, (4) *What influenced the perceptions and attitudes of GPs on health policy measures?*

Methods

Study design and setting

The PRICOV-19 study uses an online self-reported survey to conduct a multi-country cross-sectional design in 38 European countries [36]. In Germany, this study was conducted by the Institute of General Practice of the Friedrich-Alexander University Erlangen-Nürnberg (FAU).

¹ The PRICOV-19 study is a multi-country cross-sectional study researching how primary care practices were organised during the COVID-19 pandemic to guarantee safe, effective, patient-centred, and equitable care; and the pandemic's impact on the role and wellbeing of care providers. These aspects were examined through an online questionnaire.

Table 1 Main content areas and items of the questionnaire in the respective order^a

Content	Sample item
1.0 General information about the respondent and primary care	What is your position in this GP practice?
1.1 Patient flow during and before the COVID-19 pandemic:	
a. Appointments	a. Patients who made an appointment and where it is unclear whether they pose a risk of infection are called to verify this
b. Triage	b. In case the telephonic triage is performed by someone other than a GP in this GP practice and he/she needs support when assessing a call, he/she can rely on support from a GP
c. Safety management for routine primary care	c. A patient with a fever caused by an infection other than COVID-19 was seen late due to the COVID-19 protocol
1.2 Infection prevention	In this GP practice, home care nurses are currently actively contacted when their patients are diagnosed with a major infectious disease (e.g. HIV, COVID-19, hepatitis carrier status)
1.3 Information processing	In this GP practice, a fixed weekly time is provided in the agenda(s) of GPs for reviewing new guidelines or going through relevant and reliable scientific literature
1.4 Communication with patients	Does this GP practice have a brochure with information on COVID-19 to give to patients?
1.5 Dealing with health policy measure	The guidelines imposed by the government on GP practices as a consequence of COVID-19 pose a threat to the good organisation of this practice
1.6 Wellbeing	The guidelines imposed by the government on GP practices as a consequence of COVID-19 pose a threat to the personal wellbeing of the staff in this practice

^a All questions were answered with a 5-point or 7-point Likert scale using an agreement scale or could be answered with yes/no

Development, translation and validation of the questionnaire

The overall online questionnaire for all participating countries of this study was developed at Ghent University in multiple phases, including a pilot study among 159 GP practices in Flanders (Belgium). This questionnaire was developed based on a literature review and theoretical framework on quality of care [37] and patient safety culture [38]. More details are described in the study protocol (Additional File 1), including the questionnaire as an additional file [36]. The validated English questionnaire was provided to each research partner to be translated into the country's primary language. In Germany, we used a forward-backwards method to guarantee contextual conformity in translating the questionnaire into German. Two German GPs from the Institute of General Practice Erlangen independently reviewed the translated questionnaire and adapted it where necessary. Furthermore, each country could add up to ten items to the questionnaire to pursue its own research question and focus on the context of its own country, as well as primary care and health system. The amended German questions were tested for comprehensibility through cognitive interviews [39] with four GPs and two non-GPs (research associates).

Structure of the questionnaire and measures

The main questionnaire (Additional Files 2 and 3) consisted of 53 items comprising seven topics, as seen in

Table 1. Participants could also add further comments, suggestions, and feedback in an open text field. The open text field was the last item of the questionnaire, its limit was 500 words. All participants gave written informed consent on the first page of the online survey.

Furthermore, another three topics with ten items were added to the German version of the questionnaire (Additional Files 4 and 5), which mainly dealt with the context of primary care in Germany (Table 2). The focus of these particular questions lies primarily on the impact of the German health policy measures on primary care from the perspective of the GPs.

Recruitment and sample

The recruitment in Germany occurred nationwide from 01.02.2021 to 28.02.2021. The target (n) for Germany was 200 practices, following Ghent University requirements on specific ratios per country ($n=200$ for countries with more than 10,000 GPs). We assumed a response rate of 10%. The online questionnaire was directed to GPs, internists working in primary care, and GP trainees who work in single or group practices in rural, suburban or urban settings. For better readability we will refer in the following to all participants as GPs. We recruited in a snowball procedure to gain quick and broad access to the research field via anonymised e-mail dispatch lists from the general practice institutes' research practices (FAU Erlangen), other general practice institutes (throughout Germany), regional German general practices networks,

Table 2 Items added for Germany in particular^a

Content	Sample item
2.0 Implementation feasibility of standard care during COVID-19	The care of COVID (suspected) cases cannot sufficiently ensure the care of uncomplicated diseases (e.g. back pain, urinary tract infection)
2.1 Acceptance of political measures/interference of politics and society in the GP profession	The measures taken by the government concerning GP care to contain the pandemic have overwhelmed everyday practice The role of GPs has gained attention in society since the beginning of the pandemic
2.2 Structural changes in GP practice due to policy measures	The local structures of medical cooperation (e.g. interprofessional exchange, substitute organisation) have changed positively as a result of the pandemic

^a All questions were answered with a 5-point Likert scale using an agreement scale

as well as the Bavarian general practitioners' association. The participants could access the questionnaire through a link, which guided them to the online platform REDCap [40]. No incentive was given to the participants, and no reminders were sent. We sent out the survey link to 1,710 GPs. To track the data collection and the targeted n, weekly updates were sent from the REDCap server regarding the number of the already completed questionnaires. All data were collected anonymously and stored on the secure Ghent University data servers [41].

The reporting of this study is based on the *STROBE* (Strengthening the Reporting of Observational Studies in Epidemiology) recommendations [42] and the *CHERRIES* (Checklist for Reporting Results of Internet E-Surveys) checklist [43].

Data analysis

All statistical analyses were carried out with IBM SPSS Statistics 28. Boundary values of $p < 0.05$ were set for statistical significance. Data and syntax can be found in the supporting information files (Additional Files 6 and 7).

Firstly, descriptive analyses were conducted for the following topics: sample and cohort characteristics, perceiving the changes in tasks on primary care and patient treatment, perceiving the role of GPs and task shifting, perceiving the wellbeing of GPs (expanded 9-item WellbeingIndex, eWBI), as well as the assessment of the health policy measures (PolicyIndex, PI).

Secondly, two additive indices were created: The eWBI describes the frequency of distress and wellbeing among healthcare professionals. In this paper, the validated expanded 9-item Wellbeing Index version was used according to Dyrbye [44]. Cronbach's α was used to check whether all variables could be combined into an additive index (Table 6, Items: 6.0–6.8). The internal consistency was satisfying, with Cronbach's α for a positive effect of 0.76 for the eWBI. Within the eWBI, seven items had to be responded to with a yes (scored as 1) or no (scored as 0). Two items that had to be answered on a 7-point Likert scale were converted

into a new variable: response options of a scale of 1 or 2 (a low level of meaning in work) were assigned the value +1, response options of 3 to 5 (a neutral level of meaning) was given 0, and response option of 6 or 7 (a high level of meaning in work) was assigned the value -1. The 5-point Likert scale was recoded to the same scheme: those responding 'strongly disagree/disagree' having +1, those who responded 'agree/strongly agree' to having -1, and 0 for those with middle neutral responses. As in previous studies being at risk of distress is defined as a score of ≥ 2 [44]. The PI was built on the same basis as the eWBI. The PI was used to aggregate the GPs' perceptions and attitudes towards health policy measures on primary care during the pandemic. It was composed of the variables of a 5-point Likert scale (Table 8, Items: 8.0–8.3). The internal consistency was proved satisfying, with Cronbach's α for a positive effect of 0.69.

Thirdly, inferential analyses were conducted using correlation and multiple regression models. *Bravais-Pearson* [45] correlation was first calculated to see if the two indices correlate to ensure they can be used in the regression model. Regarding the multiple regression model analysis, we used the ones with the best model quality (R^2). Three models were calculated, with the metric *PI* as the dependent variable. The independent variables were introduced in three steps: (1) variables that deal with the change of role and tasks of GPs (2), the eWBI, and (3) the practice-relevant influencing factors (cohort characteristics).

Results

Sample and cohort characteristics

Three hundred forty-nine GPs filled in the questionnaire (Item 3.0) with an estimated response rate of 20.4%. At least, we had 250 fully completed questionnaires. Participants who filled out the questionnaire were mainly GPs (59.6%, Item 3.1) with mean work experience of 15 to 20 years (Item 3.2). The majority worked in single practices (62.5%, Item 3.3). Most

Table 3 German cohort characteristics

	N	%
3.0 Participants	349	100
3.1 Position/Function		
GP	208	59.6
Internists as GP	43	12.3
GP trainees	8	2.3
Missing	90	25.8
3.2 Work experience in years		
< 5	30	8.6
5—< 10	26	7.4
10—< 15	33	9.5
15—< 20	44	12.6
20—< 25	48	13.8
25—< 30	34	9.7
30—< 35	23	6.6
> 35	12	3.4
Missing	99	28.4
3.3 Practice type		
Single	218	62.5
Group	111	31.8
Missing	20	5.7
3.4 Practice location		
Rural	91	26.1
Small Town	95	27.2
Urban	76	21.8
Missing	87	24.9

practices were located in rural areas or in small towns (53.3%, Item 3.4). For further details please see Table 3.

Perceived changes in tasks on primary care and patient treatment

GPs agreed that priority was given to suspected COVID-19 cases over other patients in terms of appointments (63.3%, Item 4.0). Also, a high proportion of respondents indicated that patient care for uncomplicated cases/diseases was ensured (85%, Item 4.1). GPs experienced that the impact of the pandemic increased recommending non-COVID-19 vaccinations, e.g. influenza or pneumococcus (70.7%, Item 4.8), and requests for COVID-19 testing by asymptomatic patients (62.2%, Item 4.9). For further results, please see Table 4.

Perceived role of GPs and task shifting

Almost 40% of the GPs agreed that their role had gained more attention. However, an equal proportion also said that they did not perceive this to be the case (Item 5.0). 76% of the respondents perceived their work during the pandemic had a personal added value for them and became more meaningful to them (Item 5.1). Next, 80.3%

of them reported that their role and tasks in primary care had changed since the beginning of the pandemic (Item 5.2), with increased responsibilities (70.6%, Item 5.3). Not even half of them felt prepared for the task shifting in their professional role (49.2%, Item 5.5). 42.2% stated that they were unhappy with the task shifting in their professional role (Item 5.4). Further results can be seen in Table 5.

Perceived wellbeing of GPs (expanded 9-item WellbeingIndex, eWBI)

Half (50.5%) of the GPs stated that they felt burned out from their work (Item 6.0) and had less time for leisure time activities (38.0%, Item 6.8). 88.5% said their work during the pandemic was meaningful to them (Item 6.7). More details please see Table 6.

The risk of distress is defined with a score of ≥ 1 (*median*) according to Dyrbye [44]. Considering the eWBI, the mean is at +1.2 ($SD=2.6$), with 53.4% of the GPs situated above +1.0. This indicates that more than half of our GPs surveyed were at risk of distress due to COVID-19 during the second wave of the pandemic in Germany.

Perceived health policy measures (PolicyIndex, PI)

Around 44.5% of the respondents agreed that the government's policies and regulations on primary care practices threatened the good practice organisation (Item 7.0). Besides that, 43.3% reported that the policies threatened the personal wellbeing (Item 7.1). Furthermore, the majority agreed that no adequate government support was given (75.2%, Item 7.2), and 66.4% considered that the policies overwhelmed and negatively affected the daily life of their practice (Item 7.2). Please see Table 7.

The PI index has a range from -4.00 to +4.00 points. The lower this score, the more GPs agreed with the health policy measures; the higher this score, the less GPs agreed with the government's policies and were more dissatisfied. The *mean* was +0.7, and the *median* was +1.0. We defined being less in agreement with the policy measures as a score of ≥ 1 (*median*) according to Dyrbye [44]. Almost 60% of GPs were above the cut-off point of +1.0, meaning that there was a rather high degree of dissatisfaction among the German GPs with the health policies during the second wave of the COVID-19 pandemic.

Correlation of eWBI and PI

Table 8 shows a highly significant two-sided relationship between the eWBI and PI indices. $R=0.44$ corresponds to a medium negative effect, e.g., the higher the eWBI score (and the lower the wellbeing), the more negative the GP's health policy attitude. In the context of our previous results, this means that in the second COVID-19 wave in

Table 4 Perceived changes in tasks on primary care and patient treatment^a

	<i>n</i>	Strongly disagree	Disagree	Neutral	Agree	Strongly Agree	Mean (SD)
4.0 COVID-19 suspected cases are given priority when appointments are made.	256	6 (2.3%)	31 (12.1%)	57 (22.3%)	76 (29.7%)	86 (33.6%)	3.8 (1.1)
4.1 Care for suspected COVID-19 cases does <i>not</i> adequately ensure care for other uncomplicated cases/diseases (e.g. back pain).	259	89 (34.4%)	131 (50.6%)	22 (8.5%)	14 (5.4%)	3 (1.2%)	1.9 (0.85)
4.2 Patients with uncomplicated diseases are mainly treated by telephone consultations (e.g. back pain).	260	76 (29.2%)	127 (48.8%)	32 (12.3%)	21 (8.1%)	4 (1.5%)	2.0 (0.94)
4.3 Suspected Covid-19 cases were mainly treated via telephone consultations.	259	69 (26.6%)	81 (31.3%)	28 (10.8%)	57 (22.0%)	24 (9.3%)	2.6 (1.34)
4.4 No appointments are offered for routine consultations/examinations (e.g. back pain).	259	116 (44.8%)	103 (39.8%)	14 (5.4%)	18 (6.9%)	8 (3.1%)	1.8 (1.02)
4.5 The possibility of a telephone or video consultation relieves the burden on practice resources.	254	29 (11.4%)	68 (26.8%)	52 (20.5%)	77 (30.3%)	28 (11.0%)	3.0 (1.21)
4.6 Since the beginning of the pandemic, the practice has increasingly offered home visits to patients at risk.	259	36 (13.9%)	110 (42.5%)	48 (18.5%)	58 (22.4%)	7 (2.7%)	2.6 (1.07)
4.7 The cooperation between local practices have changed positively as a result of the pandemic (e.g. professional exchange, support).	258	47 (18.2%)	105 (40.7%)	67 (26.0%)	31 (12.0%)	8 (3.1%)	2.4 (1.02)
4.8 Protective vaccinations are increasingly recommended by the practice (e.g. influenza, pneumococcus).	259	5 (1.9%)	17 (6.6%)	54 (20.8%)	115 (44.4%)	68 (26.3%)	3.9 (0.95)
4.9 The request for COVID-19 testing by asymptomatic patients is increasing.	258	3 (1.2%)	36 (14.0%)	36 (14.0%)	103 (39.9%)	80 (31.0%)	3.9 (0.94)
4.10 COVID-19 testing <i>in</i> asymptomatic patients is more increasing as the pandemic progresses.	257	6 (2.3%)	49 (19.1%)	42 (16.1%)	100 (38.9%)	60 (23.3%)	3.6 (1.12)

^a 5-point Likert scale**Table 5** Perceived role of GPs and task shifting^a

	<i>n</i>	Strongly disagree	Disagree	Neutral	Agree	Strongly Agree	Mean (SD)
5.0 The <i>role of GPs has gained attention</i> since the beginning of the COVID-19 pandemic. ^a	255	22 (8.6%)	71 (27.8%)	61 (23.9%)	81 (31.8%)	20 (7.8%)	3.0 (1.12)
5.1 Since the COVID-19 pandemic, <i>the work I do has become more meaningful</i> to me. ^a	258	5 (1.9%)	17 (6.6%)	40 (15.5%)	107 (41.5%)	89 (34.5%)	4.0 (0.97)
5.2 <i>Primary care has changed</i> since the COVID-19 pandemic (e.g. more medical attestations, communication with health authorities). ^a	259	3 (1.2%)	17 (6.6%)	31 (12.0%)	117 (45.2%)	91 (35.1%)	4.1 (0.92)
5.3 Since the COVID-19 pandemic, <i>my responsibilities in this practice increased</i> . ^a	248	2 (0.8%)	17 (6.9%)	59 (23.8%)	87 (35.1%)	83 (35.5%)	2.9 (0.96)
5.4 I am <i>happy with the task shifting in my professional role</i> since the COVID-19 pandemic	242	18 (7.4%)	80 (34.8%)	66 (21.4%)	64 (30.6%)	14 (5.8%)	2.0 (1.04)
5.5 I <i>do not feel prepared for the task shifting in my professional role</i> since the COVID-19 pandemic. ^a	246	50 (20.3%)	71 (28.9%)	49 (18.0%)	45 (24.2%)	31 (8.5%)	1.6 (1.2)

^a 5-point Likert scale

Germany, we found that when GPs were in greater distress, they rated the health policies more negatively.

Multiple linear regression models

All three models (Table 9) were highly significant ($p < 0.001$). In the first model, a significant influence

on GPs' health policy attitudes was exerted above all by the perception of whether one felt happy with the task shifting ($\beta = -0.249$, $p < 0.001$) and whether one felt unprepared ($\beta = 0.223$, $p < 0.001$). The most substantial negative influence was dissatisfaction with the change of tasks in practice, connecting a lower satisfaction to a

Table 6 Perceived wellbeing of GPs (eWBI components)^a

	<i>n</i>	%
6.0 Have you felt burned out from your work?		
No	121	49.6
Yes	123	50.4
6.1 Have you worried that your work is hardening you emotionally?		
No	149	61.3
Yes	94	38.7
6.2 Have you often been bothered by feeling down, depressed, or hopeless?		
No	169	69.3
Yes	75	30.7
6.3 Have you fallen asleep while sitting inactive in a public place?		
No	231	95.5
Yes	11	4.5
6.4 Have you felt that all the things you had to do were piling up so high that you could not overcome them?		
No	184	76.0
Yes	58	24.0
6.5 Have you been bothered by emotional problems (such as feeling anxious, depressed, or irritable)?		
No	145	59.4
Yes	99	40.6
6.6 Has your physical health interfered with your ability to do your daily work at home and/or away from home?		
No	200	83.0
Yes	41	17.0
6.7 The work I do is meaningful to me. ^a		
1 (Strongly disagree)	-	-
2	-	-
3	2	0.8
4	7	2.9
5	19	7.8
6	80	32.9
7 (Strongly agree)	135	55.6
6.8 My work schedule leaves me enough time for my personal/family life. ^b		
1 (Strongly disagree)	26	10.7
2	66	27.3
3	59	24.4
4	63	26.0
5 (Strongly agree)	28	11.6

^a 7-point Likert scale^b 5-point Likert scale

higher PI score. In the second model, adding the factor eWBI, which had a highly significant influence on the PI ($\beta=0.347$, $p<0.001$), reduced the impact of the previously significant variables. This means that the more GPs were exposed to the risk of distress during the pandemic, the worse they assessed the government's measures. In the third model, the significance of the eWBI ($\beta=0.352$, $p<0.001$) did not change by adding the cohort characteristics, which did not influence the PI. We therefore assume that the cohort characteristics have no influence on the PI, neither directly nor indirectly mediating via

the eWBI. It can be assumed that the GPs assessed the health policy measures irrespective of their function, practice location, type of practice, or work experience. The respondents' wellbeing predominated as the most influential variable.

Discussion

Our main findings show that the German GPs experienced a considerable change in their work and practice organisation, as well as rapid changes in tasks. They felt a high responsibility within the COVID-19

Table 7 Perceived health policy measures (PI components)^a

	n	Strongly disagree	Disagree	Neutral	Agree	Strongly Agree	Mean (SD)
7.0 The guidelines imposed by the government on primary care practices as a consequence of COVID-19 pose a threat to the good organisation of this practice. ^a	246	30 (12.2%)	66 (26.8%)	55 (16.4%)	79 (38.0%)	16 (6.5%)	1.9 (1.1)
7.1 The guidelines imposed by the government on primary care practices as a consequence of COVID-19 pose a threat to the personal wellbeing. ^a	247	52 (21.1%)	24 (9.7%)	64 (25.9%)	37 (15.0%)	70 (28.3%)	1.8 (1.2)
7.2 Adequate support is provided by the government for the proper functioning of this practice. ^a	246	90 (36.6%)	95 (38.6%)	40 (16.3%)	14 (5.7%)	7 (2.8%)	1.0 (1.0)
7.3 The measures taken by the government have overburdened daily practice. ^a	254	15 (5.9%)	56 (22.0%)	55 (5.6%)	84 (39.1%)	44 (27.3%)	3.3 (1.2)

^a 5-point Likert scale

Table 8 Bravais-Pearson-Correlation between eWBI and PI

	PolicyIndex
WellbeingIndex	
Pearson-Correlation r	0.441 ^a
Sig. (two-sided)	<0.001
n	225

^a The Correlation is significant at 0.01 (two-sided)

pandemic and found that their work has become more meaningful since the pandemic. On contrary the GPs also saw a lack of political support, and many felt that they were not well prepared by the government. Furthermore, they found that the measures taken by the government have overburdened the daily practice. However, the wellbeing index showed that many respondents did not feel very strongly affected by their psychological wellbeing, but rated the implementation of the policies and especially the impact on the organisation of the practice rather negatively. The GPs' wellbeing is significantly related to how the respondents rated the German health policies: The more dissatisfied the GPs were with the task shifting, the more negative they felt about the policies decided from the government. It follows, that from the perspective of the German GPs, the health policy measures had a negative influence on everyday practice rather than being supportive. If the practice team and the GPs felt unprepared, experienced little support by the government and were dissatisfied with the changes in tasks, they evaluated the political actions more negatively. It could be deduced that if GPs were dissatisfied and overwhelmed with the measures, they may be less able to implement them. Accordingly, active inclusion of the GPs in health policy decisions

and recommendations for action and a participatory approach should be considered.

A study by Wangler et al. [46] showed that GPs, based on their own insights and experiences, do not just want advice, action and suggestions from other experts, such as policymakers. Therefore, it seems advisable to involve GPs and their experience in the planning, implementation and evaluation of measures to preparedness planning and in contingency plans [46]. Other studies [15, 21] have also highlighted the need to involve GPs in preparedness planning and showed that GPs gave low ratings to their preparedness for a pandemic. According to Siebenhofer et al. (2021), primary care is important for dealing with pandemics like COVID-19. GPs in that study said that they were confident and willing to take an active role, but they need to be provided with the appropriate ambient conditions, and information and especially be included in communicative exchange [21]. Other research [10, 47] also demonstrated that the COVID-19 pandemic had negatively affected healthcare workers in the primary ambulatory sector. Those seemed to be the most vulnerable group due to increasing responsibilities and transforming tasks. Similar findings [34, 35, 48] showed that GPs experienced a new role marked by a high-stress level combined with a deleterious work environment caused by the COVID-19 pandemic and its health policy measures. Collinset al. [34], concluded within the international comparison of the PRI-COV-19 study, that GPs with less experience, in smaller practices, and with more vulnerable patient populations had higher scores in the eWBI, thus being at a higher risk of distress. Furthermore, Groenewegen et al. (2022) found, that all over Europe, GPs and their staff members were more involved in giving information and recommendations to patients contacting the practice by phone, they were more engaged in triage of whom to refer to the hospital, and that they experienced more responsibilities [26].

Table 9 Results of linear mixed model analysis of potential predictors for the GP's perception of health policy measures (PI score, PolicyIndex) during the pandemic

Linear Mixed Models for Total PI score (PolicyIndex)					
A. Model summary					
	Correlation Coefficient r	r²	Adjusted r² p		
Modell 1	0.395	0.156	0.139 < 0.001		
Modell 2	0.505	0.255	0.237 < 0.001		
Modell 3	0.515	0.265	0.233 < 0.001		
B. Predictor coefficients					
	Regression Coefficient B	Standardised error	β	t	p
Modell 1					
(Constant)	0.193	0.767	-	0.252	0.802
Primary care changed ^a	1.203	0.633	0.127	1.900	0.059
Responsibilities increased ^a	-0.267	0.636	-0.028	-0.419	0.675
Happy with task shifting ^a	-1.197	0.316	-0.249	-3.787	< 0.001
Not feeling prepared ^a	1.015	0.299	0.223	3.394	< 0.001
Modell 2					
(Constant)	0.315	0.723	-	0.436	0.663
Primary care changed ^a	0.914	0.599	0.097	1.527	0.128
Responsibilities increased ^a	-0.535	0.601	-0.057	-0.890	0.375
Happy with task shifting ^a	-0.821	0.306	-0.170	-2.680	0.008
Not feeling prepared ^a	0.578	0.294	0.127	1.966	0.050
WellbeingIndex (eWBI)	0.312	0.060	0.347	5.222	< 0.001
Modell 3					
(Constant)	0.908	0.898	-	1.011	0.313
Primary care changed ^a	0.936	0.604	0.099	-0.868	0.123
Responsibilities increased ^a	-0.527	0.607	-0.056	-2.794	0.387
Happy with task shifting ^a	-0.863	0.309	-0.179	1.829	0.006
Not feeling prepared ^a	0.543	0.297	0.119	5.262	0.069
WellbeingIndex (eWBI)	0.317	0.060	0.352	5.262	< 0.001
Function: GP ^b	-0.341	0.369	-0.058	-0.925	0.356
Practice location: rural ^b	0.193	0.278	0.042	0.693	0.489
Type of practice: single ^b	-0.499	0.394	-0.078	-1.268	0.206
Work experience: less than 15 years ^b	0.072	0.295	0.015	0.245	0.807

^a The following 5-point Likert scale items were converted into dummies: 6.3 Primary care has changed since the COVID-19 pandemic. 6.4 Since the COVID-19 pandemic, my responsibilities in this practice increased. 6.5 I am happy with the task shifting in my professional role since the COVID-19 pandemic. 6.6 I do not feel prepared for the task shifting in my professional role since the COVID-19 pandemic

^b The following items were converted into dummies: 4.2 Position/Function. 4.3 Work experience in years. 4.4 Practice type. 4.5 practice location

Strengths and limitations

The strengths of our study are that it represents a gap in research on German primary care during the second wave of the COVID-19 pandemic in February 2021. It gives an important snapshot of the primary care practices within this time frame with a validated questionnaire and a high response rate. Furthermore, regarding our data and as it is an international cross-sectional study design, comparison with other countries becomes possible and could be an incentive for further evaluation and analysis steps.

We also see several limitations to this work. It can be assumed that the GPs who took part in the survey were rather more dissatisfied with the situation than the basic population. We therefore assume the existence of a selection bias. However, the demographic data collected does not indicate a clear major shift in the composition of the sample and the basic population. Additionally, we expect further biases, e.g. social desirability and dependence on the familiarity of the respondent with the practice organisation. Moreover, within our study, no comparison can be made between the situation before and after the pandemic and its waves.

Conclusion

The COVID-19 pandemic led to new challenges, barriers and responsibilities for German primary care. During the second wave of the pandemic, the German GPs of this study perceived their work as relevant and felt confident that they could fulfil their tasks. Nevertheless, they noticed that health policy initially hardly reacted or was delayed in reacting and supporting the outpatient sector. This led to an increased risk of distress and dissatisfaction on the part of the GPs, as an overload and high workload became apparent in the primary care setting as well, and not only at ICU level. High-quality outpatient care during the pandemic could initially only be maintained through improvisation and extreme flexibility on the part of the GPs. What was needed was a transparent pandemic plan, including the primary care sector. Consequently, it is recommended that health policy and its actors ought to take an active interest in understanding the complexity of GPs' work. They could increase their competence in relation to the primary care service, ensure that the needs of primary care are considered in preparedness plans (pandemics, climate change etc.), and include the GPs expertise concerning health policy questions.

Abbreviations

ICU	Intensive care units
PPE	Personal protective equipment
PP	Pandemic preparedness
GPs	General practitioners
PRICOV-19	Primary health care in times of COVID-19
STROBE	Strengthening the Reporting of Observational Studies in Epidemiology
CHERRIES	Checklist for Reporting Results of Internet E-Surveys
eWBI	Expanded 9-item WellbeingIndex
PI	PolicyIndex

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12875-023-02115-4>.

Additional file 1. Study protocol.

Additional file 2. Main questionnaire, English.

Additional file 3. Main questionnaire, German.

Additional file 4. Items added, English.

Additional file 5. Items added, German.

Additional file 6. Data.

Additional file 7. Syntax.

Acknowledgements

This is a secondary analysis of the PRICOV-19 study on the national level of Germany. The study was conducted in the following 37 European countries and Israel: Austria, Belgium, Bosnia and Herzegovina, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Kosovo*, Latvia, Lithuania, Luxembourg, Malta, Moldavia, North Macedonia, Norway, Poland, Portugal, Romania, Serbia, Slovenia, Spain, Sweden, Switzerland, The Netherlands, Turkey, and Ukraine; and in Israel. In total, there are more than 4,600 completed surveys. *Note: All references to Kosovo, whether the territory, institutions, or population, in this project, shall be

understood in full compliance with the United Nations Security Council Resolution 1244 and the ICJ Opinion on the Kosovo declaration of independence, without prejudice to the status of Kosovo.

The authors would like to thank all the participants who participated in this study and the whole research consortium team of the PRICOV-19 study. Additionally, a big thank you is imparted to all German Chairs of General Practice involved in the recruitment, the General Practitioners' Association, and all regional networks of GPs who assisted in distributing the questionnaire. The present work was performed in fulfilment of the requirements for obtaining the degree "Dr. rer. biol. hum." for Stefanie Stark at the Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU).

About this supplement

This article has been published as part of BMC Primary Care Volume 24 Supplement 1, 2023: COVID-19 and beyond – lessons for the future of primary care. The full contents of the supplement are available online at <https://bmcpriicare.biomedcentral.com/articles/supplements/volume-24-supplement-1>.

Authors' contributions

Conceptualisation: StSt, ES, LB. Data curation: StSt, MK, ES. Formal Analysis: StSt, FW. Methodology: StSt, SH, FW. Software: StSt. Supervision: TK, SH, FW. Writing—original draft: StSt. Writing—review & editing: ES, MK, LB, MR, EB, EvP, SW, SH, TK, FW. All authors have read and approved the final manuscript.

Funding

PRICOV-19 is set up and implemented without external funding except for a small grant from the European General Practice Research Network (EGPRN) funding to cover the data cleaning; no grant number applies. The German part of the study was financed from budgetary funds which had no influence on the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Availability of data and materials

All relevant data is available at BMC online and within this paper as Supporting Information files.

Declarations

Ethics approval and consent to participate

The Research Ethics Committee of Ghent University Hospital approved the protocol of the PRICOV-19 study (BC-07617) [36]. The Ethics Committee of the Friedrich-Alexander University Erlangen-Nürnberg approved the overall study, the German sub-study, and the data collection (379_20 B). All data is anonymised, and all raw data that could lead to the identification of the respondents was permanently removed.

Consent for publication

Not applicable.

Competing interests

None declared.

Author details

¹Institute of General Practice, Friedrich-Alexander University Erlangen-Nürnberg (FAU), Erlangen, Germany. ²Department of Public Health and Primary Care, Ghent University, Ghent, Belgium. ³Department for Sociology, University of Education, Schwäbisch Gmünd, Germany. ⁴General Practice, Faculty of Medicine, University of Augsburg, Augsburg, Germany.

Received: 16 December 2022 Accepted: 18 July 2023

Published online: 11 October 2023

References

- Balog, A. *Soziale Phänomene: Identität, Aufbau und Erklärung*. Springer-Verlag, 2007.
- Kunin M, Engelhard D, Piterman L, et al. Response of general practitioners to infectious disease public health crises: an integrative systematic review of the literature. *Disaster Med Public Health Prep.* 2013;7(5):522–33.

3. de Sutter A, Llor C, Maier M, et al. Family medicine in times of 'COVID-19': A generalists' voice. *Eur J Gen Pract.* 2020;26(1):58–60.
4. Kumar R, Nedungalaparambil NM, Mohanan N. Emergency and primary care collaboration during COVID-19 pandemic: a quick systematic review of reviews. *J Fam Med Prim Care.* 2020;9(8):3856.
5. Alexander JC. *What Makes a Social Crisis?: The Societalization of Social Problems.* Wiley: 2019.
6. Corona Schneider A. In the crisis and after the crisis. *Sozialwirtschaft aktuell.* 2020;30(8):1–3.
7. Krumpal I. *Sociology in Times of a Pandemic.* Work Paper Institut Social Univ Leipzig. 2020;79:1–16.
8. Kaldewey D. What does systemic relevance mean in times of pandemic? *Berl J Soziol.* 2022;32(1):7–33.
9. Schilling J, Tolksdorf K, Marquis A, et al. The different phases of the COVID-19 pandemic in Germany: a descriptive analysis from January 2020 to February 2021. *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz.* 2021;64(9):1093–106.
10. Cochrane Effective Practice and Organisation of Care Group, Pollock A, Campbell P, Cheyne J, et al. Interventions to support the resilience and mental health of frontline health and social care professionals during and after a disease outbreak, epidemic or pandemic: a mixed methods systematic review. *Cochrane Database Syst Rev.* 2020;11(CD013779):4–28.
11. Provenzano DA, Sitzman BT, Fiorentino SA, et al. Clinical and economic strategies in outpatient medical care during the COVID-19 pandemic. *Reg Anesth Pain Med.* 2020;45(8):579–85.
12. Rawaf S, Allen LN, Stigler FL, et al. Lessons on the COVID-19 pandemic, for and by primary care professionals worldwide. *European Journal of General Practice.* 2020;26(1):129–33.
13. Qiu H, Tong Z, Ma P, Hu M, Peng Z, Wu W, Du B. China Critical Care Clinical Trials Group (CCCCTG). Intensive care during the coronavirus epidemic. *Intensive Care Med.* 2020;46(4):576–8.
14. Stang ASM, Jöckel K-H. Estimated use of intensive care beds due to COVID-19 in Germany over time. *Dtsch Arztebl Int.* 2020;117(19):329–35.
15. Stöcker A, Demirer I, Gunkel S, et al. Stockpiled personal protective equipment and knowledge of pandemic plans as predictors of perceived pandemic preparedness among German general practitioners. *PLoS ONE.* 2021;16(8):e0255986.
16. Gassen A. *Pandemie-Management in der ambulanten Versorgung: Analyse des bisherigen Verlaufs – Strategien und Maßnahmen für die Zukunft.* Medizinisches Wissenschaftliche Verlagsgesellschaft. 2020. Available at: https://www.kbv.de/media/sp/Gassen_Pandemie-Management_amb_Versorgung_WCFM_MWV_2020.pdf. Accessed 04 Sept 2023.
17. Webb E, Hernández-Quevedo C, Williams G, et al. Providing health services effectively during the first wave of COVID-19: a cross-country comparison on planning services, managing cases, and maintaining essential services. *Health Policy.* 2022;126(5):382–90.
18. Winkelmann J, Webb E, Williams GA, et al. European countries' responses in ensuring sufficient physical infrastructure and workforce capacity during the first COVID-19 wave. *Health Policy.* 2022;126(5):362–72.
19. Renaa T, Brekke M. GPs' experiences of the COVID-19 pandemic - a focus group study. *Tidsskrift for Den norske legeforening.* 2022;142(12):1–7.
20. Krist AH, DeVoe JE, Cheng A, et al. Redesigning primary care to address the COVID-19 pandemic in the midst of the pandemic. *Ann Fam Med.* 2020;18(4):349–54.
21. Siebenhofer A, Huter S, Avian A, et al. COVI-Prim survey: challenges for Austrian and German general practitioners during initial phase of COVID-19. *PLoS One.* 2021;16(6):e0251736.
22. Scherr A. Corona Crisis. *Sozial Extra.* 2020;44:172–6.
23. Paffenholz P, Peine A, Hellmich M, et al. Perception of the 2020 SARS-CoV-2 pandemic among medical professionals in Germany: results from a nationwide online survey. *Emerg Microbes Infect.* 2020;9(1):1590–9.
24. [fos/aerzteblatt.de: Corona crisis: Care physician can order cooperation in Bavaria. Deutsches Ärzteblatt.](https://www.aerzteblatt.de/nachrichten/111496/Coronakrise-Versorgungsarzt-kann-in-Bayern-Mitarbeiter-anordnen) 2020. Available at: <https://www.aerzteblatt.de/nachrichten/111496/Coronakrise-Versorgungsarzt-kann-in-Bayern-Mitarbeiter-anordnen>. Accessed 04 Sept 2023.
25. Schreyögg J. Corona crisis meets structural problems in health care. *Wirtschaftsdienst, Springer.* 2020;100(4):226–7.
26. Groenewegen P, van Poel E, Spreuwenberg P, et al. Has the COVID-19 pandemic led to changes in the tasks of the primary care workforce? An International Survey among General Practices in 38 Countries (PRICOV-19). *Int J Environ Res Public Health.* 2022;19(22):15329.
27. Sarti TD, Lazarini, WS, Fontenelle LF, et al. What is the role of primary health care in the COVID-19 pandemic? *Epidemiologia e serviços de saúde.* 2020;29.
28. WHO. *Primary health care.* 2020.
29. Dunlop C, Howe A, Li D, et al. The coronavirus outbreak: the central role of primary care in emergency preparedness and response. *BJGP Open.* 2020;4(1):bjgpopen20X101041.
30. Kearon J, Risdon C. The role of primary care in a pandemic: reflections during the COVID-19 pandemic in Canada. *J Prim Care Community Health.* 2020;11:2150132720962871.
31. Greenfield G, Foley K, Majeed A. Rethinking primary care's gatekeeper role. *BMJ.* 2016;354:i4803.
32. Kurotschka PK, Serafini A, Demontis M, et al. General practitioners' experiences during the first phase of the COVID-19 pandemic in Italy: a Critical Incident Technique study. *Front Public Health.* 2021;9:19.
33. Gray R, Sanders C. A reflection on the impact of COVID-19 on primary care in the United Kingdom. *J Interprof Care.* 2020;34(5):672–8.
34. Collins C, Clays E, van Poel E, et al. Distress and wellbeing among general practitioners in 33 countries during COVID-19: results from the cross-sectional PRICOV-19 study to inform health system interventions. *Int J Environ Res Public Health.* 2022;19(9):5675.
35. Jefferson L, Golder S, et al. GP wellbeing during the COVID-19 pandemic: a systematic review. *Br J Gen Pract.* 2022;72(718):e325–33.
36. Van Poel E, VandenBussche P, Klemenc-Ketis Z, Willems S. How did general practices organize care during the COVID-19 pandemic: the protocol of the cross-sectional PRICOV-19 study in 38 countries. *BMC Fam Pract.* 2022;23:11.
37. IoM. *Improving the 21st century health care system. Crossing the Quality Chasm: A New Healthcare System for the 21st Century.* 2001.
38. Dovey SM, Meyers DS, Phillips RL, et al. A pre-liminary taxonomy of medical errors in family practice. *BMJ Qual Saf.* 2002;11(3):233–8.
39. Collins D. Pretesting survey instruments: an overview of cognitive methods. *Qual Life Res.* 2003;12:229–38.
40. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform.* 2009;42(2):377–81.
41. Van Poel E, Vanden Bussche P, Klemenc-Ketis Z, Willems S. How did general practices organize care during the COVID-19 pandemic: The protocol of the cross-sectional PRICOV-19 study in 38 countries. *BMC Prim Care.* 2022;23(1):1–11.
42. von Elm E, Altman DG, et al. Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ.* 2007;335(7624):806–8.
43. Eysenbach G. Improving the quality of Web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res.* 2004;6(3):e34.
44. Dyrbye LN, Satele D, Shanafelt T. Ability of a 9-item well-being index to identify distress and stratify quality of life in US workers. *J Occup Environ Med.* 2016;58:810–7.
45. Artusi R, Verderio P, Marubini E. Bravais-Pearson and Spearman correlation coefficients: meaning, test of hypothesis and confidence interval. *Int J Biol Markers.* 2002;17(2):148–51.
46. Wangler J, Jansky M. Approaches to securing primary care. *ZfA Zietschrf für Allgemeinmedizin.* 2022;98(7–8):250–6.
47. Serrano-Ripoll MJ, Meneses-Echavez JF, Ricci-Cabello I, et al. Impact of viral epidemic outbreaks on mental health of healthcare workers: A rapid systematic review and meta-analysis. *J Affect Disord.* 2020;277:347–57.
48. Astill Wright L, Gnanapragasam S, Downes AJ. Managing COVID-19 related distress in primary care: principles of assessment and management. *BMC Fam Pract.* 2021;22(1):73.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.